

Asymmetric Dark Matter Stability from Continuous Flavor Symmetries

Fady Bishara

Fermilab & University of Cincinnati

Fermilab
September 11th, 2014



Based on: FB and J. Zupan. arXiv:1408.3852

- ▷ **Motivation**
- ▷ ADM, DM stability, and flavor
- ▷ Asymmetric Dark Matter (ADM) mass
- ▷ ADM lifetime
- ▷ Mediator models
- ▷ Experimental constraints

Outline

- ▶ Motivation
- ▶ ADM, DM stability, and flavor
- ▶ Asymmetric Dark Matter (ADM) mass
- ▶ ADM lifetime
- ▶ Mediator models
- ▶ Experimental constraints

Outline

- ▶ Motivation
- ▶ ADM, DM stability, and flavor
- ▶ Asymmetric Dark Matter (ADM) mass
- ▶ ADM lifetime
- ▶ Mediator models
- ▶ Experimental constraints

Outline

- ▶ Motivation
- ▶ ADM, DM stability, and flavor
- ▶ Asymmetric Dark Matter (ADM) mass
- ▶ ADM lifetime
- ▶ Mediator models
- ▶ Experimental constraints

Outline

- ▷ Motivation
- ▷ ADM, DM stability, and flavor
- ▷ Asymmetric Dark Matter (ADM) mass
- ▷ ADM lifetime
- ▷ Mediator models
- ▷ Experimental constraints

Outline

- ▷ Motivation
- ▷ ADM, DM stability, and flavor
- ▷ Asymmetric Dark Matter (ADM) mass
- ▷ ADM lifetime
- ▷ Mediator models
- ▷ Experimental constraints

Motivation

- ▶ There is overwhelming evidence for the existence of DM yet the SM model lacks a candidate
- ▶ There is a coincidence $\Omega_\chi/\Omega_B = 5.4$; could there be a link?
- ▶ We expect New Physics (NP) at the TeV scale to address the hierarchy problem
- ▶ However, NP cannot have generic flavor structure
 - Large FCNCs if $\Lambda_{\text{NP}} \sim \text{TeV}$ (NP flavor problem)
- ▶ Can flavor suppression lead to DM stability?

ADM, DM stability and flavor

There is a vast literature on the topic. Some examples include

▷ ADM

Hooper, March-Russell & West [hep-ph/0410114], Kaplan, Luty & Zurek [aXv:0901.4117], Feldstein & Fitzpatrick [aXv:1003.5662], Dutta & Kumar [aXv:1012.1341], Cohen, Phalen, Pierce & Zurek [aXv:1005.1655], Falkowski, Ruderman & Volansky [aXv:1101.4936]

▷ MFV

Kamenik & Zupan [aXv:1107.0623], Batell, Pradler & Spannowsky [aXv:1105.1781], Batell, Lin & Wang [aXv:1309.4462], SUSY MFV: Csaki, Grossman & Heidenreich [aXv:1111.1239], Monteux & Cornell [aXv:1404.5952]

▷ Lepton and quark flavored DM

Agrawal, Blanchet, Chacko & Kilic [aXv:1109.3516], Kumar & Tulin [aXv:1303.0332], Agrawal, Batell, Hooper & Lin [aXv:1404.1373]

▷ Beyond MFV

Agrawal, Blanke & Gemmler [aXv:1405.6709]

The roadmap

- ▶ Flavor & SM gauge singlet DM charged under $U(1)_{(B-L)}$
 \Rightarrow DM is either a Dirac fermion or a complex scalar
- ▶ Assume that $B \neq 0$ and $L = 0$ to focus the discussion
- ▶ DM is a color singlet \Rightarrow carries integer Baryon number
- ▶ Will not assume any discrete symmetry to stabilize DM

Goal

A cosmologically stable DM with $\Lambda_{\text{NP}} \sim \mathcal{O}(\text{TeV})$

The roadmap

- ▶ Flavor & SM gauge singlet DM charged under $U(1)_{(B-L)}$
 \Rightarrow DM is either a Dirac fermion or a complex scalar
- ▶ Assume that $B \neq 0$ and $L = 0$ to focus the discussion
- ▶ DM is a color singlet \Rightarrow carries integer Baryon number
- ▶ Will not assume any discrete symmetry to stabilize DM

Goal

A cosmologically stable DM with $\Lambda_{\text{NP}} \sim \mathcal{O}(\text{TeV})$

ADM mass

Assumptions

- ▶ $B - L$ is a conserved quantum number
- ▶ Symmetric component efficiently annihilated


In this case, the ADM mass (with SM field content) is given by¹

$$m_\chi = m_p \frac{\Omega_\chi}{\Omega_B} \left(\frac{B}{B-L} \right) \left(\frac{B-L}{\Delta_\chi} \right) = (12.5 \pm 0.8 \text{ GeV}) \frac{1}{(B-L)_\chi^{\text{sum}}}$$

where $\Delta_\chi \equiv (n_\chi - \bar{n}_\chi)/s$ and $(B-L)_\chi^{\text{sum}} \equiv \sum_i \hat{g}_\chi^i (B-L)_\chi^i$.

¹Harvey & Turner, Phys.Rev. D42 (1990) 3344-3349; Feldstein & Fitzpatrick, arXiv:1003.5662.

ADM mass in the presence of New Physics (NP)



Asymmetric EFT operators

The lowest dimensional asymmetric operators are of the form

$$\mathcal{L} = \sum_i \frac{\mathcal{C}_i}{\Lambda^{(D_i-4)}} \chi \mathcal{O}_i^{\text{SM}},$$

$$\text{with}^1 \mathcal{O}^{\text{SM}} = [u^c]^{n_u} [d^c]^{n_d} [q^*]^{n_q},$$

$$\text{and } \begin{cases} (n_d + n_u + n_q) \bmod 3 = 0 \\ n_d - n_u - n_q/2 = 0 \end{cases}$$

¹The fields u^c and d^c are the $SU(2)_L$ singlet up and down type quark fields while q is the $SU(2)_L$ doublet quark field in two component spinor notation.

Freeze-out temperature of asymm. operators

- ▶ The NDA estimate for the freeze-out temperature is (FN flavor breaking)

$$T_f \sim \left(1.66 \times \sqrt{g_*} (16\pi^2)^3 \frac{8\pi}{C^2} \frac{\Lambda^{12}}{M_{\text{pl}}} \right)^{1/11} \simeq 480 \text{ GeV} \left(\frac{\Lambda}{1.9 \text{ TeV}} \right)^{12/11}$$

- ▶ The EFT scale $\Lambda > 1.9 \text{ TeV}$ is bounded by indirect detection searches.
- ▶ Dominated by the $2 \rightarrow 5$ process.
- ▶ DM number is conserved below T_f .



Metastability and flavor breaking

To calculate the DM lifetime we must

- ▶ Choose the flavor structure. We will consider two flavor breaking scenarios: Minimal Flavor Violation (MFV) and Froggatt-Nielsen (FN)
- ▶ Rotate to the mass eigenbasis. We will work in the down mass basis where

$$u^c \rightarrow u_{\text{MASS}}^c, \quad d^c \rightarrow d_{\text{MASS}}^c, \quad q = \begin{pmatrix} u \\ d \end{pmatrix} \rightarrow \begin{pmatrix} V_{\text{CKM}} u_{\text{MASS}} \\ d_{\text{MASS}} \end{pmatrix}.$$

and the Yukawa matrices are

$$Y_D \rightarrow Y_D^{\text{diag}}, \quad Y_U \rightarrow V_{\text{CKM}} Y_U^{\text{diag}}$$

- ▶ Using Naive dimensional analysis (NDA), estimate DM total width

Metastability and flavor breaking

To calculate the DM lifetime we must

- ▶ Choose the flavor structure. We will consider two flavor breaking scenarios: Minimal Flavor Violation (MFV) and Froggatt-Nielsen (FN)
- ▶ Rotate to the mass eigenbasis. We will work in the down mass basis where

$$u^c \rightarrow u_{\text{MASS}}^c, \quad d^c \rightarrow d_{\text{MASS}}^c, \quad q = \begin{pmatrix} u \\ d \end{pmatrix} \rightarrow \begin{pmatrix} V_{\text{CKM}} u_{\text{MASS}} \\ d_{\text{MASS}} \end{pmatrix}.$$

and the Yukawa matrices are

$$Y_D \rightarrow Y_D^{\text{diag}}, \quad Y_U \rightarrow V_{\text{CKM}} Y_U^{\text{diag}}$$

- ▶ Using Naive dimensional analysis (NDA), estimate DM total width

Metastability and flavor breaking

To calculate the DM lifetime we must

- ▶ Choose the flavor structure. We will consider two flavor breaking scenarios: Minimal Flavor Violation (MFV) and Froggatt-Nielsen (FN)
- ▶ Rotate to the mass eigenbasis. We will work in the down mass basis where

$$u^c \rightarrow u_{\text{MASS}}^c, \quad d^c \rightarrow d_{\text{MASS}}^c, \quad q = \begin{pmatrix} u \\ d \end{pmatrix} \rightarrow \begin{pmatrix} V_{\text{CKM}} u_{\text{MASS}} \\ d_{\text{MASS}} \end{pmatrix}.$$

and the Yukawa matrices are

$$Y_D \rightarrow Y_D^{\text{diag}}, \quad Y_U \rightarrow V_{\text{CKM}} Y_U^{\text{diag}}$$

- ▶ Using Naive dimensional analysis (NDA), estimate DM total width

Metastability and flavor breaking

To calculate the DM lifetime we must

- ▶ Choose the flavor structure. We will consider two flavor breaking scenarios: Minimal Flavor Violation (MFV) and Froggatt-Nielsen (FN)
- ▶ Rotate to the mass eigenbasis. We will work in the down mass basis where

$$u^c \rightarrow u_{\text{MASS}}^c, \quad d^c \rightarrow d_{\text{MASS}}^c, \quad q = \begin{pmatrix} u \\ d \end{pmatrix} \rightarrow \begin{pmatrix} V_{\text{CKM}} u_{\text{MASS}} \\ d_{\text{MASS}} \end{pmatrix}.$$

and the Yukawa matrices are

$$Y_D \rightarrow Y_D^{\text{diag}}, \quad Y_U \rightarrow V_{\text{CKM}} Y_U^{\text{diag}}$$

- ▶ Using Naive dimensional analysis (NDA), estimate DM total width

Metastability and flavor breaking

To calculate the DM lifetime we must

- ▶ Choose the flavor structure. We will consider two flavor breaking scenarios: Minimal Flavor Violation (MFV) and Froggatt-Nielsen (FN)
- ▶ Rotate to the mass eigenbasis. We will work in the down mass basis where

$$u^c \rightarrow u_{\text{MASS}}^c, \quad d^c \rightarrow d_{\text{MASS}}^c, \quad q = \begin{pmatrix} u \\ d \end{pmatrix} \rightarrow \begin{pmatrix} V_{\text{CKM}} u_{\text{MASS}} \\ d_{\text{MASS}} \end{pmatrix}.$$

and the Yukawa matrices are

$$Y_D \rightarrow Y_D^{\text{diag}}, \quad Y_U \rightarrow V_{\text{CKM}} Y_U^{\text{diag}}$$

- ▶ Using Naive dimensional analysis (NDA), estimate DM total width

Minimal Flavor Violation¹ (MFV)

- ▶ \mathcal{L}_{SM} enjoys an enhanced symmetry G_F in the limit $m_q \rightarrow 0$
- ▶ $G_F = SU(3)_Q \times SU(3)_U \times SU(3)_D$
- ▶ Symmetry is retained if Yukawa matrices are promoted to spurions that transform under G_F as

$$Y_U \sim (\mathbf{3}, \bar{\mathbf{3}}, \mathbf{1}), \quad Y_D \sim (\mathbf{3}, \mathbf{1}, \bar{\mathbf{3}})$$

- ▶ The Yukawa interactions $u^c Y_U^\dagger q H$, $d^c Y_D^\dagger q H^c$ are then formally invariant under G_F

The SM Yukawas are the only source of flavor breaking.

¹D'Ambrosio, Giudice, Isidori & Strumia [hep-ph/0207036]

Minimal Flavor Violation¹ (MFV)

- ▶ \mathcal{L}_{SM} enjoys an enhanced symmetry G_F in the limit $m_q \rightarrow 0$
- ▶ $G_F = SU(3)_Q \times SU(3)_U \times SU(3)_D$
- ▶ Symmetry is retained if Yukawa matrices are promoted to spurions that transform under G_F as

$$Y_U \sim (\mathbf{3}, \bar{\mathbf{3}}, \mathbf{1}), \quad Y_D \sim (\mathbf{3}, \mathbf{1}, \bar{\mathbf{3}})$$

- ▶ The Yukawa interactions $u^c Y_U^\dagger q H$, $d^c Y_D^\dagger q H^c$ are then formally invariant under G_F

The SM Yukawas are the only source of flavor breaking.

¹D'Ambrosio, Giudice, Isidori & Strumia [hep-ph/0207036]

Example: $B = 1$ operators with MFV

$$\mathcal{O}_1^{(B=1)} = (\chi u^c)(d^c d^c), \quad \mathcal{O}_2^{(B=1)} = (\chi q_\rho^*)(d^c q_\sigma^*)\epsilon^{\rho\sigma}$$

$$\begin{aligned} \mathcal{O}_1^{(B=1)} &= (\chi u_\alpha^c Y_U^\dagger Y_D)_K (d_{N\beta}^c d_{M\gamma}^c) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma} \\ &\rightarrow (\chi u_{\text{MASS}}^c Y_U^{\text{diag}\dagger} V_{\text{CKM}}^\dagger Y_D^{\text{diag}})_{K\alpha} ([d_{\text{MASS}}^c]_{N\beta} [d_{\text{MASS}}^c]_{M\gamma}) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma}, \end{aligned}$$

$$\begin{aligned} \mathcal{O}_2^{(B=1)} &= (\chi q_{K\alpha i}^*) ([d_\beta^c Y_D^\dagger]_N q_{M\gamma j}^*) \epsilon^{ij} \epsilon^{KNM} \epsilon^{\alpha\beta\gamma} \\ &\rightarrow (\chi u_{\text{MASS}}^* V_{\text{CKM}}^\dagger)_{K\alpha} ([d_{\text{MASS}}^c Y_D^{\text{diag}\dagger}]_{N\beta} [d_{\text{MASS}}^*]_{M\gamma}) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma}, \end{aligned}$$

$$\begin{aligned} \Gamma_\chi^{(1)} &\sim \frac{(y_t y_b)^2}{8\pi} \left(\frac{m_\chi}{\Lambda}\right)^4 \left(\frac{1}{16\pi^2} \frac{m_t \Lambda_{\text{QCD}}}{m_W^2}\right)^2 \frac{m_\chi}{16\pi^2} \\ &= 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{5.3 \cdot 10^6 \text{TeV}}{\Lambda}\right)^4, \end{aligned}$$

$$\Gamma_\chi^{(2)} \sim \frac{|y_b V_{ub}|^2}{8\pi} \left(\frac{m_\chi}{\Lambda}\right)^4 \frac{m_\chi}{16\pi^2} = 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{4.8 \cdot 10^7 \text{TeV}}{\Lambda}\right)^4$$

Example: $B = 1$ operators with MFV

$$\mathcal{O}_1^{(B=1)} = (\chi u^c)(d^c d^c), \quad \mathcal{O}_2^{(B=1)} = (\chi q_\rho^*)(d^c q_\sigma^*)\epsilon^{\rho\sigma}$$

$$\begin{aligned} \mathcal{O}_1^{(B=1)} &= (\chi u_\alpha^c Y_U^\dagger Y_D)_{K\alpha} (d_{N\beta}^c d_{M\gamma}^c) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma} \\ &\rightarrow (\chi u_{\text{MASS}}^c Y_U^{\text{diag}\dagger} V_{\text{CKM}}^\dagger Y_D^{\text{diag}})_{K\alpha} ([d_{\text{MASS}}^c]_{N\beta} [d_{\text{MASS}}^c]_{M\gamma}) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma}, \\ \mathcal{O}_2^{(B=1)} &= (\chi q_{K\alpha i}^*) ([d_\beta^c Y_D^\dagger]_{N\gamma} q_{M\gamma j}^*) \epsilon^{ij} \epsilon^{KNM} \epsilon^{\alpha\beta\gamma} \\ &\rightarrow (\chi u_{\text{MASS}}^* V_{\text{CKM}}^\dagger)_{K\alpha} ([d_{\text{MASS}}^c Y_D^{\text{diag}\dagger}]_{N\beta} [d_{\text{MASS}}^*]_{M\gamma}) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma}, \end{aligned}$$

$$\begin{aligned} \Gamma_\chi^{(1)} &\sim \frac{(y_t y_b)^2}{8\pi} \left(\frac{m_\chi}{\Lambda}\right)^4 \left(\frac{1}{16\pi^2} \frac{m_t \Lambda_{\text{QCD}}}{m_W^2}\right)^2 \frac{m_\chi}{16\pi^2} \\ &= 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{5.3 \cdot 10^6 \text{TeV}}{\Lambda}\right)^4, \\ \Gamma_\chi^{(2)} &\sim \frac{|y_b V_{ub}|^2}{8\pi} \left(\frac{m_\chi}{\Lambda}\right)^4 \frac{m_\chi}{16\pi^2} = 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{4.8 \cdot 10^7 \text{TeV}}{\Lambda}\right)^4 \end{aligned}$$

Example: $B = 1$ operators with MFV

$$\mathcal{O}_1^{(B=1)} = (\chi u^c)(d^c d^c), \quad \mathcal{O}_2^{(B=1)} = (\chi q_\rho^*)(d^c q_\sigma^*)\epsilon^{\rho\sigma}$$


$$\begin{aligned} \mathcal{O}_1^{(B=1)} &= (\chi u_\alpha^c Y_U^\dagger Y_D)_K (d_{N\beta}^c d_{M\gamma}^c) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma} \\ &\rightarrow (\chi u_{\text{MASS}}^c Y_U^{\text{diag}\dagger} V_{\text{CKM}}^\dagger Y_D^{\text{diag}})_{K\alpha} ([d_{\text{MASS}}^c]_{N\beta} [d_{\text{MASS}}^c]_{M\gamma}) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma}, \end{aligned}$$

$$\begin{aligned} \mathcal{O}_2^{(B=1)} &= (\chi q_{K\alpha i}^*) ([d_\beta^c Y_D^\dagger]_N q_{M\gamma j}^*) \epsilon^{ij} \epsilon^{KNM} \epsilon^{\alpha\beta\gamma} \\ &\rightarrow (\chi u_{\text{MASS}}^* V_{\text{CKM}}^\dagger)_{K\alpha} ([d_{\text{MASS}}^c Y_D^{\text{diag}\dagger}]_{N\beta} [d_{\text{MASS}}^*]_{M\gamma}) \epsilon^{KNM} \epsilon^{\alpha\beta\gamma}, \end{aligned}$$

$$\begin{aligned} \Gamma_\chi^{(1)} &\sim \frac{(y_t y_b)^2}{8\pi} \left(\frac{m_\chi}{\Lambda}\right)^4 \left(\frac{1}{16\pi^2} \frac{m_t \Lambda_{\text{QCD}}}{m_W^2}\right)^2 \frac{m_\chi}{16\pi^2} \\ &= 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{5.3 \cdot 10^6 \text{TeV}}{\Lambda}\right)^4, \end{aligned}$$

$$\Gamma_\chi^{(2)} \sim \frac{|y_b V_{ub}|^2}{8\pi} \left(\frac{m_\chi}{\Lambda}\right)^4 \frac{m_\chi}{16\pi^2} = 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{4.8 \cdot 10^7 \text{TeV}}{\Lambda}\right)^4$$

Example: $B = 1$ operators with MFV




$$\Gamma_{\chi}^{(1)} \sim \frac{(y_t y_b)^2}{8\pi} \left(\frac{m_{\chi}}{\Lambda}\right)^4 \left(\frac{1}{16\pi^2} \frac{m_t \Lambda_{\text{QCD}}}{m_W^2}\right)^2 \frac{m_{\chi}}{16\pi^2}$$

$$= 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{5.3 \cdot 10^6 \text{TeV}}{\Lambda}\right)^4,$$

$$\Gamma_{\chi}^{(2)} \sim \frac{|y_b V_{ub}|^2}{8\pi} \left(\frac{m_{\chi}}{\Lambda}\right)^4 \frac{m_{\chi}}{16\pi^2} = 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024}\right)^2 \left(\frac{4.8 \cdot 10^7 \text{TeV}}{\Lambda}\right)^4$$

Example: $B = 1$ operators with MFV



$$\Gamma_{\chi}^{(1)} \sim \frac{(y_t y_b)^2}{8\pi} \left(\frac{m_{\chi}}{\Lambda} \right)^4 \left(\frac{1}{16\pi^2} \frac{m_t \Lambda_{\text{QCD}}}{m_W^2} \right)^2 \frac{m_{\chi}}{16\pi^2}$$

$$= 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024} \right)^2 \left(\frac{5.3 \cdot 10^6 \text{TeV}}{\Lambda} \right)^4,$$

$$\Gamma_{\chi}^{(2)} \sim \frac{|y_b V_{ub}|^2}{8\pi} \left(\frac{m_{\chi}}{\Lambda} \right)^4 \frac{m_{\chi}}{16\pi^2} = 6.6 \cdot 10^{-51} \text{GeV} \left(\frac{y_b}{0.024} \right)^2 \left(\frac{4.8 \cdot 10^7 \text{TeV}}{\Lambda} \right)^4$$



DM leading decays and EFT scale

B	ADM model			MFV		FN		
	Dim.	m_χ [GeV]	decay	τ [s]	Λ [TeV]	decay	τ [s]	Λ [TeV]
1	6	6.2	$\chi \rightarrow bus$	10^{26}	4.0×10^6	$\chi \rightarrow bus$	10^{26}	8.1×10^8
2	10	3.1	$\chi \rightarrow udsuds$	10^{26}	0.63	$\chi \rightarrow udsuds$	10^{26}	2.5
3	15	2.1	forbidden	∞	–	forbidden	∞	–

Table: Leading decay modes for the $B = \{1, 2, 3\}$ operators with MFV and FN flavor breaking. The scale Λ_* is calculated such that the lifetime of the DM $\tau \sim 10^{26}$ [s]. The decay of ADM with $B = 3$ is kinematically forbidden.

DM leading decays and EFT scale

ADM model				MFV		FN		
B	Dim.	m_χ [GeV]	decay	τ [s]	Λ [TeV]	decay	τ [s]	Λ [TeV]
1	6	6.2	$\chi \rightarrow bus$	10^{26}	4.0×10^6	$\chi \rightarrow bus$	10^{26}	8.1×10^8
2	10	3.1	$\chi \rightarrow udsuds$	10^{26}	0.63	$\chi \rightarrow udsuds$	10^{26}	2.5
3	15	2.1	forbidden	∞	–	forbidden	∞	–


Table: Leading decay modes for the $B = \{1, 2, 3\}$ operators with MFV and FN flavor breaking. The scale Λ_* is calculated such that the lifetime of the DM $\tau \sim 10^{26}$ [s]. The decay of ADM with $B = 3$ is kinematically forbidden.

DM leading decays and EFT scale

B	ADM model			MFV		Anarchic		
	Dim.	m_χ [GeV]	decay	τ [s]	Λ [TeV]	decay	τ [s]	Λ [TeV]
1	6	6.2	$\chi \rightarrow bus$	10^{26}	4.0×10^6			
2	10	3.1	$\chi \rightarrow udsuds$	10^{26}	0.63	$\chi \rightarrow udd udd$	10^{26}	7.3
3	15	2.1	forbidden	∞	—			

Table: Leading decay modes for the $B = \{1, 2, 3\}$ operators with MFV and FN flavor breaking. The scale Λ_* is calculated such that the lifetime of the DM $\tau \sim 10^{26}$ [s]. The decay of ADM with $B = 3$ is kinematically forbidden.

ADM lifetime



Ackermann et al. [aXv:1205.6474]; Ibarra, Lamperstorfer, & Silk [aXv:1309.2570]; Aguilar et al. [Phys.Rev.Lett. 110, 141102 (2013)]; Covi, Grefe, Ibarra, & Tran [aXv:0912.3521]; Desai et al. [aXv:hep-ex/0404025]; Zhao & Zurek [aXv:1401.7664]

Mediator models

MFV model with scalar mediators

$$\begin{aligned}\mathcal{L}_{\text{INT}} \supset & \kappa_1 [\phi_L]_{\gamma}^{AB} (q_{A,\alpha i}^* q_{B,\beta j}^*) \epsilon^{ij} \epsilon^{\alpha\beta\gamma} + \kappa_2 [\varphi_L]_A^{\alpha\beta} (q_{B,\alpha i}^* q_{C,\beta j}^*) \epsilon^{ij} \epsilon^{ABC} \\ & + \kappa_3 [Y_D]_X^A [\phi_R]_{A,\alpha} (d_{Y,\beta}^c d_{Z,\gamma}^c) \epsilon^{\alpha\beta\gamma} \epsilon^{XYZ} + \kappa_4 \chi^\dagger [\phi_L]_{\alpha}^{AB} [\varphi_L]_A^{\alpha\beta} [\phi_R]_{B,\beta} \\ & + h.c.\end{aligned}$$

The gauge and global charge assignment for the three scalar mediators, ϕ_L , φ_L and ϕ_R , in the first UV completion toy model for which we also assume the MFV flavor breaking pattern

Field	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	G_F	$U(1)_{B-L}$
ϕ_L	$\bar{\mathbf{3}}$	$\mathbf{1}$	$1/3$	$(\mathbf{6}, \mathbf{1}, \mathbf{1})$	$2/3$
φ_L	$\mathbf{6}$	$\mathbf{1}$	$1/3$	$(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1})$	$2/3$
ϕ_R	$\bar{\mathbf{3}}$	$\mathbf{1}$	$-2/3$	$(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1})$	$2/3$

MFV model with scalar mediators

$$\begin{aligned}\mathcal{L}_{\text{INT}} \supset & \kappa_1 [\phi_L]_\gamma^{AB} (q_{A,\alpha i}^* q_{B,\beta j}^*) \epsilon^{ij} \epsilon^{\alpha\beta\gamma} + \kappa_2 [\varphi_L]_A^{\alpha\beta} (q_{B,\alpha i}^* q_{C,\beta j}^*) \epsilon^{ij} \epsilon^{ABC} \\ & + \kappa_3 [Y_D]_X^A [\phi_R]_{A,\alpha} (d_{Y,\beta}^c d_{Z,\gamma}^c) \epsilon^{\alpha\beta\gamma} \epsilon^{XYZ} + \kappa_4 \chi^\dagger [\phi_L]_\alpha^{AB} [\varphi_L]_A^{\alpha\beta} [\phi_R]_{B,\beta} \\ & + h.c.\end{aligned}$$

The gauge and global charge assignment for the three scalar mediators, ϕ_L , φ_L and ϕ_R , in the first UV completion toy model for which we also assume the MFV flavor breaking pattern

Field	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	G_F	$U(1)_{B-L}$
ϕ_L	$\bar{\mathbf{3}}$	$\mathbf{1}$	$1/3$	$(\mathbf{6}, \mathbf{1}, \mathbf{1})$	$2/3$
φ_L	$\mathbf{6}$	$\mathbf{1}$	$1/3$	$(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1})$	$2/3$
ϕ_R	$\bar{\mathbf{3}}$	$\mathbf{1}$	$-2/3$	$(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1})$	$2/3$

MFV model with scalar mediators


$$\begin{aligned}\mathcal{L}_{\text{INT}} \supset & \kappa_1 [\phi_L]_\gamma^{AB} (q_{A,\alpha i}^* q_{B,\beta j}^*) \epsilon^{ij} \epsilon^{\alpha\beta\gamma} + \kappa_2 [\varphi_L]_A^{\alpha\beta} (q_{B,\alpha i}^* q_{C,\beta j}^*) \epsilon^{ij} \epsilon^{ABC} \\ & + \kappa_3 [Y_D]_X^A [\phi_R]_{A,\alpha} (d_{Y,\beta}^c d_{Z,\gamma}^c) \epsilon^{\alpha\beta\gamma} \epsilon^{XYZ} + \kappa_4 \chi^\dagger [\phi_L]_\alpha^{AB} [\varphi_L]_A^{\alpha\beta} [\phi_R]_{B,\beta} \\ & + h.c.\end{aligned}$$

The gauge and global charge assignment for the three scalar mediators, ϕ_L , φ_L and ϕ_R , in the first UV completion toy model for which we also assume the MFV flavor breaking pattern

Field	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	G_F	$U(1)_{B-L}$
ϕ_L	$\bar{\mathbf{3}}$	$\mathbf{1}$	$1/3$	$(\mathbf{6}, \mathbf{1}, \mathbf{1})$	$2/3$
φ_L	$\mathbf{6}$	$\mathbf{1}$	$1/3$	$(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1})$	$2/3$
ϕ_R	$\bar{\mathbf{3}}$	$\mathbf{1}$	$-2/3$	$(\bar{\mathbf{3}}, \mathbf{1}, \mathbf{1})$	$2/3$

MFV model with scalar mediators

$$\begin{aligned} \mathcal{L}_{\text{INT}} \supset & \kappa_1 [\phi_L]_\gamma^{AB} (q_{A,\alpha i}^* q_{B,\beta j}^*) \epsilon^{ij} \epsilon^{\alpha\beta\gamma} + \kappa_2 [\varphi_L]_A^{\alpha\beta} (q_{B,\alpha i}^* q_{C,\beta j}^*) \epsilon^{ij} \epsilon^{ABC} \\ & + \kappa_3 [Y_D]_X^A [\phi_R]_{A,\alpha} (d_{Y,\beta}^c d_{Z,\gamma}^c) \epsilon^{\alpha\beta\gamma} \epsilon^{XYZ} + \kappa_4 \chi^\dagger [\phi_L]_\alpha^{AB} [\varphi_L]_A^{\alpha\beta} [\phi_R]_{B,\beta} \\ & + h.c. \end{aligned}$$



FN model with scalar and fermionic mediators


$$\mathcal{L}_{\text{INT}} \supset g_{q,AB} \phi_\gamma \left(q_{A,\alpha i}^{*j} q_{B,\beta j}^{*k} \right) \epsilon^{ij} \epsilon^{\alpha\beta\gamma} + g_{d,A} \phi^{*\alpha} \left(d_{A,\alpha}^c \psi \right) + g_\chi \chi (\psi^c \psi^c) + h.c.$$

Gauge and $B - L$ charges of the mediators ϕ and ψ in the second UV model. We also assume FN flavor breaking pattern

Field	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$U(1)_{B-L}$
ϕ	$\bar{\mathbf{3}}$	$\mathbf{1}$	$1/3$	$2/3$
ψ	$\mathbf{1}$	$\mathbf{1}$	0	1


FN model with scalar and fermionic mediators

$$\mathcal{L}_{\text{INT}} \supset g_{q,AB} \phi_\gamma \left(q_{A,\alpha i}^{*j} q_{B,\beta j}^{*k} \right) \epsilon^{ij} \epsilon^{\alpha\beta\gamma} + g_{d,A} \phi^{*\alpha} \left(d_{A,\alpha}^c \psi \right) + g_\chi \chi (\psi^c \psi^c) + h.c.$$




Flavor constraints

Mediators contribute to $\Delta_F = 2$ processes at the one loop level via



Flavor constraints

Mediators contribute to $\Delta_F = 2$ processes at the one loop level via








As in the SM, there is a GIM cancellation in these diagrams and the contribution is additionally suppressed by the internal quark Yukawa.

Flavor constraints



	MFV		FN	
	$\kappa_{1,2} <$	$m_{\phi_L, \varphi_L} >$	$g_{q,d} <$	$m_\phi >$
$K^0 - \bar{K}^0$	0.33	2.9 TeV	0.63	570 GeV
$B_d - \bar{B}_d$	1.3	710 GeV	0.54	1 TeV
$B_s - \bar{B}_s$	1.3	780 GeV	0.59	840 GeV
$D^0 - \bar{D}^0$	30	34 GeV	4.3	56 GeV

Table: The 95 % C.L. bounds on the MFV and FN mediator models from meson mixing. Taking $m_{\phi_L} = m_{\varphi_L} = m_\phi = 1\text{TeV}$ and $\kappa_1 = \kappa_2$ gives the upper bounds on the couplings in the 2nd column and 4th column for $g_q = g_d$. Taking $\kappa_{1,2} = g_{q,d} = 1$ gives lower bounds on the mediator masses in the 3rd and 5th columns. The mass of the fermion in the FN model is fixed to $m_\psi = 20\text{ GeV}$.


Flavor constraints – MFV mediator model




Flavor constraints – FN mediator model



Collider signatures: single and pair production




Collider signatures: paired dijets constraints



Search for New Physics in the Paired Dijet Mass Spectrum – CMS. [arXiv:1302.0531]

Collider signatures: $2b$ jets + MET




► The NDA decay length of ψ is given by

$$\begin{aligned} c\tau(\psi \rightarrow bbc) &\sim \left(g_q^2 g_d^2 \lambda^8 \frac{1}{8\pi} \frac{1}{16\pi^2} \frac{m_\psi^5}{m_\phi^4} \right)^{-1} \\ &\sim 30\text{m} \left(\frac{20\text{ GeV}}{m_\psi} \right)^5 \left(\frac{m_\phi}{750\text{ GeV}} \right)^4 \left(\frac{0.03}{g_q g_d} \right)^2 \end{aligned}$$

Collider signatures: $2b$ jets + MET


Constraints from sbottom pair production



Search for direct production of a pair of bottom squarks – CMS. [PAS-SUS-13-018]

Collider signatures: $2b$ jets + MET

Constraints from sbottom pair production



Search for direct production of a pair of bottom squarks – CMS. [PAS-SUS-13-018]

Summary & conclusions

- ▶ Showed that flavor symmetries can allow us to have a cosmologically stable ADM even if the DM is not charged under the flavor group
- ▶ The mediators between the visible and dark sectors can be at the TeV scale without giving rise to dangerous FCNCs
- ▶ The mediator models can have interesting signatures at the LHC

Backup

$U(1)$ Froggatt-Nielsen¹ (FN) model

- ▷ Spontaneously broken horizontal $U(1)$ symmetry
- ▷ Quarks carry horizontal charges under this $U(1)$
- ▷ E.g., horizontal charge assignment that gives phenomenologically satisfactory quark masses and CKM matrix elements²

$$H(q, d^c, u^c) \Rightarrow \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} q \\ d^c \\ u^c \end{matrix} & \begin{pmatrix} 3 & 2 & 0 \\ 3 & 2 & 2 \\ 3 & 1 & 0 \end{pmatrix} \end{matrix}$$

- ▷ Wilson coefficients $\mathcal{C} = \lambda^{|\sum_i H_i|}$, where $\lambda = 0.2$

¹Froggatt & Nielsen [Nucl.Phys. B147 (1979) 277]

²Leurer, Nir & Seiberg [hep-ph/9310320], [hep-ph/9212278]